

Production and Fungicidal Activity Assesment of Wood- waste Liquid Smoke

by Budi Rahmat

Submission date: 08-Mar-2021 01:43AM (UTC-0800)

Submission ID: 1527286530

File name: and_Fungicidal_Activity_Assesment_of_Wood-waste_Liquid_Smoke.pdf (580.85K)

Word count: 3626

Character count: 18991



PRODUCTION AND FUNGICIDAL ACTIVITY ASSESMENT OF WOOD-WASTE LIQUID SMOKE



Budy Rahmat ^{*1}✉, Apip Hermawan ¹, Dedi Natawijaya ¹, Endang Surahman ²


20

¹ Department of Agrotechnology, Siliwangi University, Tasikmalaya, Indonesia

² Department of Physical Education, Siliwangi University, Tasikmalaya, Indonesia

17

DOI: <https://doi.org/10.29121/granthaalayah.v8.i10.2020.1970>



Article Type: Research Article

Article Citation: Budy Rahmat, Apip Hermawan, Dedi Natawijaya, and Endang Surahman. (2020). PRODUCTION AND FUNGICIDAL ACTIVITY ASSESMENT OF WOOD-WASTE LIQUID SMOKE. International Journal of Research - GRANTHAALAYAH, 8(10), 285-291. <https://doi.org/10.29121/granthaalayah.v8.i10.2020.1970>

Received Date: 11 October 2020

Accepted Date: 31 October 2020

Keywords:

Concentration
Fungicide
Liquid Smoke
Soybean
Wood Waste

ABSTRACT

Liquid smoke is known to contain compounds that can control plant disease pathogens. This study aims to produce wood-waste liquid smoke and determine its effectiveness as a fungicide on plant pathogens. This research was conducted in two experimental stages, namely: (i) in vitro test as a preliminary test of the effectiveness of teak waste liquid smoke at concentrations of 0, 0.5, 1, 1.5, 2, and 2.5%; and (ii) in vivo test was arranged in randomized block design consisting of seven levels of liquid smoke concentration, namely 0, 1, 2, 3, 4, 5, and 6%, each of which was repeated four times. The results showed that the pyrolysis of 1 kg of wood waste was produced with the proportions of liquid smoke, charcoal and tar, respectively: 312 mL, 31 g, 367 g and the uncondensed gases. Treatment of liquid smoke in the in vivo test showed that a concentration of 1 to 2.5% liquid smoke was able to suppress the growth of the pathogenic fungus *Sclerotium rolfsii* 100%. The treatment of liquid smoke in the in vivo test showed an effect on inhibition of the growth diameter of fungal colonies, suppressing the disease occurrence, and suppressing the lesion diameter.

1. INTRODUCTION

Sustainable agricultural development requires environmentally friendly pest control. In practice, vegetable pesticides are used, one of which is liquid smoke. Internal studies regarding the use of liquid smoke have been carried out and have the potential as a plant regulator for papaya [1], then as a larvacide and repellent for *Spodoptera litura* [2], and able to control the main pests of rice namely the golden snail [3]. As an antifungal, it affects the strawberry fruit rot pathogens [4].

Liquid smoke is known to contain compounds that can control disease-causing pathogens in plants. The compounds in liquid smoke that affect the anti-fungal properties of liquid smoke are phenols and acids.

Phenols and acids are produced by pyrolysis of lignin and hemicellulose. The by-product of processed teak wood has the same properties both in the structure and chemical properties of teak. Therefore, teak wood shavings have the potential to be turned into liquid smoke [5].

Liquid smoke from charcoal furnace processing was able to inhibit the growth of soil-borne fungi *Sclerotium oryzae*, *Pythium spp.*, and *Rhizoctonia solani* with the results showing that liquid smoke was able to inhibit pathogenic

4

activity from a concentration of 2% [6]. Volatile compounds and non-volatile compounds in liquid smoke can inhibit the growth of the fungal mycelium *Rhizoctonia solani* and *Sclerotinia sclerotiorum* [7].

This study aims to produce liquid smoke and its effectiveness as a fungicide.

2. MATERIALS AND METHODS

2.1. MATERIAL

The tools used in this experiment were a set of liquid smoke pyrolysis and distillation tools, container bottles, measuring cups, test tubes, beaker glasses, erlenmeyers, ose needles, cork borer, gauze, aluminum foil, petri dishes, spatulas, tweezers, micropipettes, electric scale, bunsen, gas stove, oven, laminar air flow, thermohigrometer, autoclave, plastic tray, plastic box, hand sprayer, calipers, funnel, filter paper, pH meter, glass object, microscope.

This experiment used materials including: teak wood shavings liquid-smoke, Potato Dextrose Agar (PDA) media, aquades, *Sclerotium rolfsii* isolate, 3% NaOCl, 70% alcohol, 1% FeCl₃, soil, soybean seeds, manure.

2.2. EXPERIMENTAL SETUP

This experiment was arranged in a randomized block design, which consisted of 7 levels of treatment and was repeated 4 times. The treatment tested was the concentration of teak wood shavings, namely: K₀ = control (aquades), K₁ = 1%, K₂ = 2%, K₃ = 3%, K₄ = 4%, K₅ = 5%, and K₆ = 6%.

The data were analyzed by using analysis of variance (Anova) and continued with the Duncan's multiple different test [8].

2.3. PROCEDURE OF EXPERIMENT

Making liquid smoke uses the following tools and procedures used by Rahmat et al. [3] and Rahmat et al. [5]. Raw liquid smoke is redistilled once to produce liquid smoke in grade 2.

In vitro testing was carried out by mixing liquid smoke with PDA media according to the desired concentration. The concentrations of liquid smoke to be used are 0, 0.5, 1, 1.5, 2, and 2.5%. The next step, the *Sclerotium rolfsii* culture was cut using a cork borer with a diameter of 5 mm, then placed right in the center of the media, done in duplicate, and observed for 4 days.

In vivo testing was carried out on 5 day old soybean plants. The plants were then sterilized using 3% NaOCl by soaking them for 5 minutes then rinsing with distilled water and drying them. The plastic box was filled with seven plants which had been surface sterilized and maintained their physiological activity. The plant roots were wrapped in cotton soaked in water. Pathogen inoculation by attaching pieces of hyphae with a diameter of 5 mm to each plant stem and then incubated for 6 hours. After that, the application of liquid smoke was carried out by soaking the inoculated plants in a liquid smoke solution according to the treatment for 5 minutes. Then the soybean plants are returned to the box in their original position.

2.4. DATA COLLECTION

The parameters observed included: characteristics of teak wood shavings liquid smoke, identification of pathogenic isolates, in-vitro anti-fungal index, disease incidence rate and length of lesions. The characteristics of liquid smoke observed included color, aroma, pH, specific gravity, acid content and phenol test results. Fungi identification was done by observing the characteristics and morphology of hyphae and sclerotia on PDA media both macroscopically and microscopically.

In the *in vitro* test, the fungisidal action (FA) value was obtained using the following formula:

$$FA = 1 - \frac{CT}{CC} \times 100\%$$

Note: CT = colony diameter in treatment; CC = colony diameter in control

The disease occurrence rate (DO) was calculated by the following equation:

$$DO = \frac{SI}{ST} \times 100\%$$

Note: SI = number of infected seedling; ST = number of total observed seedling

From the disease occurrence rate (DO), the suppression of disease occurrence (SDO) can be determined with the following equation:

$$SDO = \left(\frac{DOC - DOT}{DOC} \right) \times 100\%$$

Note: DOC = disease occurrence rate in control; DOT = disease occurrence rate in treatment

3. RESULTS AND DISCUSSIONS

24

3.1. CHARACTERISTICS OF LIQUID SMOKE

The characteristics of teak wood shavings liquid smoke produced are: reddish-brown in color, pH 3, and a very strong smoke aroma. The yield resulting from the pyrolysis process of 1 kg of teak wood shavings was produced with the proportions of liquid smoke, charcoal and tar, respectively: 312 mL, 31 g, 367 g and the uncondensed gases. The lower yield of liquid smoke than that produced by Rahmat et al. [4] and Wagini and Sukaryono [7] possibly caused by differences in the moisture content of the materials, pyrolysis kiln design, temperature and pyrolysis time.

The results of redistillation obtained that the average proportion of liquid smoke was 82.25%, and 17.75% tar and uncondensable compounds. Redistilled liquid smoke changes color, pH and aroma. Grade 2 liquid smoke has characteristics: pale yellow color, pH 2, and reduced smoke aroma. This teak wood shavings liquid smoke distillate has a density of 1.0107, the acid content is 7.026%. The results of the characterization of liquid smoke can be seen in Table 1, with a comparison of the standard Japanese liquid smoke distillate [9].

The phenol test results show positive, there was a dark black color, which indicates the presence of phenol content in the liquid smoke.

Table 1: Comparison of sample liquid smoke and Japanese quality standard

Parameter	Liquid smoke distillate	Liquid smoke by Japanese standards
pH	2	1,3 to 3,7
Density	1.0107	> 1,001
Color	Pale yellow	Colorless to reddish brown
Transparency	Transparent	Not cloudy (transparent)
Acid content	7.026%	1 to 18%

31

3.2. FUNGICIDAL ACTION OF LIQUID SMOKE IN VITRO

In the control treatment, the pathogenic fungal mycelium *S. rolfsii* began to grow on agar media at 9 hours after incubation. At 24 hours after incubation, the diameter of the mycelium colonies reached 1.1 cm. Then at 4 days after incubation the fungal mycelium growth has filled the entire area of the petri dish (Fig.1.). The maximum growth of *S. rolfsii* on PDA media is in the range of 4 and 7 days after incubation. Furthermore, the measurement of the fungicidal action rate the liquid smoke treatment was only carried out for four days [10]. The results of measuring the fungicidal action can be seen in Table 2.

Table 2: Effect of liquid smoke concentration on fungicidal activity

Liquid Smoke Concentration	Fungicidal activity rate (%)			
	1 DAI	2 DAI	3 DAI	4 DAI
0,5%	100	73,469	66,667	46,389
1%	100	100	100	100

29

1,5%	100	100	100	100
2%	100	100	100	100
2,5%	100	100	100	100

Note: DAI = days after incubation

10

In media with a liquid smoke concentration of 0.5% the fungi began to grow at 2 days after incubation. The decrease in antifungal activity began at 2, 3 and 4 days after incubation. However, at a concentration of 1 to 2.5% there was no decrease in antifungal activity, it remained 100%, until the measurement time was complete.




Figure 1: The top row is a medium with a concentration of 1% liquid smoke (K_1) at 1 to 3 DAI; and the bottom row is a medium with a concentration of 0 % liquid smoke (K_0 /control) at 1 to 3 DAI.

The fungisidal action shows the ability of liquid smoke to inhibit the development of fungi, namely the presence of phenol compounds in liquid smoke. The mechanism of activity of phenolic compounds as antimicrobial substances includes reactions in cell membranes that cause increased permeability and result in loss of cell contents, inactivation of essential enzymes, and destruction or functional inactivation of genetic material. The higher the phenol concentration will precipitate all cell proteins, while the lower the concentration will effectively inhibit essential enzymes [11].

Organic acids, such as acetic acid, propionic acid, lactic acid, benzoic acid, and salicylic acid can act as antimicrobials mainly due to the formation of free H^+ ions. At concentrations of acetic acid above 0.5% it is able to penetrate into cells and denature cell plasma proteins. Propionic acid can inhibit microbes by blocking the cellular metabolic system through inhibition of enzyme activity. In addition, propionic acid is able to reduce the intracellular pH value which can result in inhibition or cell killing [12].

3.3. DISEASE OCCURRENCE RATE *IN VIVO*

2 Table 3 showed that, since 1 day after application there was a difference in the occurrence of disease between the control and all treatments from the concentration of liquid smoke. The incidence in the control was 70.36%, while in all treatments the concentration of liquid smoke was able to withstand 0%. This shows that in the liquid smoke there was anti-fungal activity of liquid smoke that suppressed *S. rolfsii* growth.

There was no disease occurrence since 1 day after application in all liquid smoke treatments, because the active compounds in liquid smoke were at an intolerance level for fungal life. At 2 days after application, it began to show differences in the disease occurrence, namely the decrease in the occurrence with the increasing concentration of phenol and organic acids in the liquid smoke solution.


Table 3: Effect of liquid smoke concentration on disease occurrence rate

Liquid smoke concentration	Disease occurrence rate (%)							
	1 DAI		2 DAI		3 DAI		4 DAI	
K ₀ (0%)	70,36	a	100,00	a	100,00	a	100	a
K ₁ (1%)	0,00	b	32,50	b	42,50	b	47,50	b
K ₂ (2%)	0,00	b	9,17	c	21,67	c	32,50	b
K ₃ (3%)	0,00	b	4,17	c	16,07	c	25,24	bc
K ₄ (4%)	0,00	b	3,57	c	10,71	c	10,71	cd
K ₅ (5%)	0,00	b	0,00	c	0,00	d	4,17	d
K ₆ (6%)	0,00	b	0,00	c	0,00	d	3,57	d

Note: Numbers followed by the same letter indicates not significantly different according to Duncan's Multiple range test level of 5%. The data was transformed by $\sqrt{(x + 0.5)}$. DAI = days after inoculation.

The facts mentioned above show that liquid smoke is most effective as an antifungal in affecting the incidence rate of disease due to *Sclerotium rolfsii* at a concentration of 4 to 6%. At these concentrations, the quantity of phenolic compounds is thought to be able to inhibit the development of fungi. The difference in the concentration of the liquid smoke given causes a difference in the quantity of the compounds present in the liquid smoke. The higher the concentration of liquid smoke, the higher the anti-fungal activity. Accordingly the study that reported the higher concentration of liquid smoke that was applied was able to reduce the percentage of strawberries infected with the *Rhizopus stolonifer* fungus [4]. Furthermore, other study reported that at a concentration of 5% coconut shell liquid smoke was able to inhibit the growth of *Fusarium oxysporum* and *Colletotrichum gloeosporioides* [13].

The role of phenol compounds and their derivatives is associated with impaired integrity and permeability of fungal cell membranes which can cause permanent damage to cell walls and membranes. Observations using a Scanning Electron Microscope (SEM) regarding the effect of eugenol on *Zygosaccharomyces rouxii* with the results showing that eugenol at the minimum inhibitory concentration (0.4 μl / ml) is able to cause the cell surface to shrink, while increasing the concentration to the minimum fungicide concentration (0.8 μl / ml) causes cell destruction and the presence of a torn cell surface [14].

**Figure 2:** *S. rolfsii* attack on K₀ (control); and on K₁ (1% concentration of liquid smoke).

3.4. DIAMETER OF THE LESION *IN VIVO*

The results showed that there was an effect of anti-fungal activity on the length of fungal lesions infecting plant stems. At 1 day after application, there was a very significant effect of the antifungal activity of liquid smoke, and at

2 days after application, there was an increase in the diameter of the lesion in the control treatment, followed by the concentration of liquid smoke from 1 to 4%. The highest increase in lesion length occurred in the control and it was different for all the liquid smoke treatments that were tried. In the same treatment, the concentration of liquid smoke there was a tendency that the more the concentration of liquid smoke would decrease the length of the fungal lesion. This is in line with the study results showed that coconut shell liquid smoke in the range of 5 to 10% was able to suppress the development of pod rot disease by pressing the length of spots on the fruit caused by *Phytophthora palmivora* [15].

Table 4: Effect of liquid smoke concentration on diameter of lesion

Liquid smoke concentration	Diameter of the lesion (mm)							
	1 DAI		2 DAI		3 DAI		4 DAI	
K ₀ (0%)	4,61	a	11,14	a	16,53	a	22,58	a
K ₁ (1%)	0,00	b	2,01	b	4,09	b	4,52	b
K ₂ (2%)	0,00	b	0,77	bc	1,69	c	1,83	c
K ₃ (3%)	0,00	b	0,40	c	1,09	cd	1,18	cd
K ₄ (4%)	0,00	b	0,03	c	0,80	cd	0,86	cd
K ₅ (5%)	0,00	b	0,00	c	0,00	d	0,20	d
K ₆ (6%)	0,00	b	0,00	c	0,00	d	0,16	d

Note: Numbers followed by the same letter indicates not significantly different according to Duncan's Multiple range test level of 5%. The data was transformed by $\sqrt{x + 0.5}$. DAI = days after inoculation.

The process of formation of lesions through several series of biochemical processes with enzymatic mechanisms. The mechanism of enzyme inactivation was thought to be the most influencing factor in the formation of lesions on plant stems. Phenolic compounds as antimicrobials in several mechanisms that occurred include inactivation of essential enzymes, as well as functional destruction or inactivation of genetic material in the fungal body [13]. This was supported by the study, which found that the mechanism of phenolic compounds as anti-fungal was through inhibition of enzyme activity needed when infecting plants [16].

The effect of acid content in low concentrations of liquid smoke was thought to have little effect on fungal development, because fungi are able to adapt to fairly acidic environmental conditions. However, environmental conditions such as low pH are thought to directly acidify the cytoplasm, damage the surface tension of the membrane and lose the active transport of food through the membrane, causing the destabilization of various functions and structures of cell components [13].

4. CONCLUSION

The results of the study provided the following conclusions:

- 1) The pyrolysis of 1 kg of teak wood shavings was produced with the proportions of liquid smoke, charcoal and tar, respectively: 312 mL, 31 g, 367 g and the uncondensed gases.
- 2) Liquid smoke from teak wood shavings was effective as fungicide on the damping off pathogen (*Sclerotium rolfsii*). The effectiveness of this liquid smoke includes: inhibition of the diameter of fungal colonies, suppression of disease occurrence rates, and suppression of lesion diameter.
- 3) Concentrations of liquid smoke ranging from 4% were effective as a fungicide to control *Sclerotium rolfsii* as a pathogen for soybean damping off.

9

SOURCES OF FUNDING

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST

The author have declared that no competing interests exist.

International Journal of Research -GRANTHAALAYAH

13

ACKNOWLEDGMENT

The authors are grateful for the support of the Directorate General of Research and Development of the Ministry of Research, Technology and the Higher Education Republic of Indonesia under the Grant Agreement of the Basic Research Number 209 /SP2H/LT/DRPM/2019.

REFERENCES

- [1] Rahmat, B., Pangesti, D., Natawijaya, D., Sufyadi, D.. Generating Wood-waste Vinegar and Its Effectiveness as Plant Growth Regulator and Insect Pest Repellent, *BioResources Journal*, 9, 2014, 6350-6360.
- [2] Rahmat, B., Kurniati, F., Hartini, E..Mahogany wood-waste vinegar as larvacide on *Spodoptera litura*, *oResources Journal*, 10, 2015, 6741-6750.
- [3] Rahmat, B., Kurniati, F., Pajar, L. The effectiveness of teak wood-sawdust liquid smoke and areca-nut extract as a pesticide on *Pomacea canaliculata*, *American Journal of Agricultural and Biological Sciences*, 14, 2019, 69-74.
- [4] Setiawan, W., Effect of coconut shell liquid smoke on soft rot pathogens (*Rhizopus stolonifer*) in strawberries (*Fragaria x ananassa*), Thesis. Agrotechnology Study Program, Faculty of Agriculture, Siliwangi University, Tasikmalaya. 2015
- [5] Wagini, R., and Sukaryono, S. Physical characteristics and content of teak wood powder liquid smoke as a source of biofuel energy, *Indonesian Journal of Physics*, 8, 2009, 1-15.
- [6] Chalermisan, Y. and Peerapan, S. Wood vinegar: by-product from rural charcoal kiln and its role in plant protect, *Asian Journal of Food and Agro-Industry*, 2009, 189-195.
- [7] Saberi, Sarpeleh A., Askary, H., and Rafiei, F. The effectiveness of wood vinegar in controlling *Rhizoctonia solani* and *Sclerotinia sclerotiorum* in green house-cucumber, *Internasional Journal of Agriculture and Natural Resources*, 1, 2013, 039-043.
- [8] Gomez, K. and Gomez, A.A. Statistical Procedures for Agricultural Research, John Wiley & Sons. Inc., 1984.
- [9] Montazeri, N., Oliveira, A.C.M., Himelbloom, B.H., Leigh, M.B., and Crapo, C.A. Chemical characterization of commercial liquid smoke products, *Food Science & Nutrition* 1, 2013,102-115.
- [10] Dwetha, G.S.. Studies on variability and management of *Sclerotium rolfsii* Sacc. causing wilt of stevia". Thesis. Department of Plant Pathology College of Agriculture, Dharwad University of Agricultural Sciences, Dharwad. 2012.
- [11] Maqsood, S., Benjakul, S., and Shahidi, F. Emerging Role of Phenolic Compounds as Natural Food Additives in Meat and Fish Products, *Critical Reviews in Food Science and Nutrition*, 53, 2013, 162-179.
- [12] Luck, E., and Jager, M. Antimicrobial Food Additives, Characteristics, Uses and Effects, Springer-Verlag Berlin Heidelberg GmbH. Pp. 257. 1995
- [13] Aisyah, I., Juli, N. and G. Pari, G. Utilization of coconut shell liquid smoke to control fungi that cause anthracnose disease and fusarium wilt in cucumbers, *Journal of Forest Products Research*, 31,2013,170-178..
- [14] Cai, R., Hu, M., Zhang, Y., Niu, C., Yue, T., Yuan, Y. and Wang, Z. Antifungal activity and mechanism of citral, limonene and eugenol against *Zygosaccharomyces rouxii*" *LWT-Food Science and Technology*. 2019.
- [15] Yunita, I. Siswanto., And Sharbino. Effect of coconut shell liquid smoke on *P. palmivora* which causes pod rot disease in cocoa. *Journal of Plantation and Tropical Land*, 8, 2018, 91-97.
- [16] Oramahi, H.A., Rusmiyanto, E., Wardoyo, P. and Kustiati.. "Efficacy of liquid smoke from bengkirai wood against *Phytophthora citrophthora*". *Indonesian Journal of Plant Protection*, 22, 2018: 160-166.

Production and Fungicidal Activity Assessment of Wood-waste Liquid Smoke

ORIGINALITY REPORT

18%

SIMILARITY INDEX

14%

INTERNET SOURCES

15%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1

docplayer.net

Internet Source

1%

2

www.myfoodresearch.com

Internet Source

1%

3

www.ncsu.edu

Internet Source

1%

4

journals.plos.org

Internet Source

1%

5

seminar.uad.ac.id

Internet Source

1%

6

scipub.org

Internet Source

1%

7

krishikosh.egranth.ac.in

Internet Source

1%

8

uclouvain.be

Internet Source

1%

9

Denise Howel. "Trends in the prevalence of

obesity and overweight in English adults by age and birth cohort, 1991–2006", Public Health Nutrition, 2010

Publication

1%

10

M. Faisal, Asri Gani, Husni. "UTILIZATION OF LIQUID SMOKE FROM OIL PALM KERNEL SHELL TO PRESERVE MACKEREL", Rasayan Journal of Chemistry, 2018

Publication

1%

11

www.intechopen.com

Internet Source

1%

12

e.bangor.ac.uk

Internet Source

1%

13

N. Sree Rajani, V. Bhargavi Reddy. "UNDERSTANDING YOGASUTRA: A FRAMEWORK FOR ECOLOGICAL VIRTUE ETHICS", International Journal of Research - GRANTHAALAYAH, 2017

Publication

1%

14

worldwidescience.org

Internet Source

1%

15

Rizki Wuri Widaryuni, Oktira Roka Aji. "In vitro antifungal activity of lime peels extract (Citrus aurantifolia (Cristm.) Swingle.) against Fusarium oxysporum Schlecht. em. Snyder & Hans.", Symposium of Biology Education (Symbion),

1%

-
- | | | |
|----|--|-----|
| 16 | Submitted to Universitas Warmadewa
Student Paper | 1 % |
|----|--|-----|
-
- | | | |
|----|--|------|
| 17 | www.researchgate.net
Internet Source | <1 % |
|----|--|------|
-
- | | | |
|----|---|------|
| 18 | A Indriati, D D Hidayat, D A Darmajana.
"Performance Test And Economic Evaluation Of
Pyrolysis Furnace Small Scale-Worth The
Business For Producing Liquid Smoke", IOP
Conference Series: Earth and Environmental
Science, 2019
Publication | <1 % |
|----|---|------|
-
- | | | |
|----|--|------|
| 19 | www.ejournalofscience.org
Internet Source | <1 % |
|----|--|------|
-
- | | | |
|----|---|------|
| 20 | N Ratnaningsih, R R El Akbar, E Hidayat.
"Effect of chronotype and student learning time
on mathematical ability based on self-regulated
learning", Journal of Physics: Conference
Series, 2018
Publication | <1 % |
|----|---|------|
-
- | | | |
|----|--|------|
| 21 | uygarbilgi.wordpress.com
Internet Source | <1 % |
|----|--|------|
-
- | | | |
|----|--|------|
| 22 | pure.uva.nl
Internet Source | <1 % |
|----|--|------|
-

23	www.freepatentsonline.com Internet Source	<1 %
24	Satrijo Saloko, Purnama Darmadji, Bambang Setiaji, Yudi Pranoto. "Antioxidative and antimicrobial activities of liquid smoke nanocapsules using chitosan and maltodextrin and its application on tuna fish preservation", Food Bioscience, 2014 Publication	<1 %
25	research.library.mun.ca Internet Source	<1 %
26	H A Dien, R I Montolalu, S Berhimpon. "Liquid smoke inhibits growth of pathogenic and histamine forming bacteria on skipjack fillets", IOP Conference Series: Earth and Environmental Science, 2019 Publication	<1 %
27	sigc.edu Internet Source	<1 %
28	www.ajas.info Internet Source	<1 %
29	Montazeri Mahmood, Hiyam Hosseini, Behroz Jokar. "COMPARATIVE STUDY OF COMPLETED SUICIDE BASED ON SEASON AND DISEASES IN BUSHEHR PROVINCE, (2006-2015)", International Journal of Research	<1 %

30

U. I. Meko Ayub, Berhimpon Siegfried, Ketut Suwetja I, G. Ijong Frans. "Moisture sorption isotherm of Tuna Sei, smoked with liquid smoke", International Journal of Fisheries and Aquaculture, 2016

Publication

<1 %

31

D A Susbandi, F Swastawati, S Suharto. " Flavor characteristics and quality of mackerel () "otak-otak" affected by various liquid smoke addition methods ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

32

Fronthea Swastawati, Ambaryanto, Bambang Cahyono, Ima Wijayanti, Diana Chilmawati. " Characterizations of milkfish () meatballs as effect of nanoencapsulation liquid smoke addition ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

Off