

# A QUALITATIVE ANALYSIS OF TANNIN TYPE AND TANNIN CONTENT IN MENIRAN TEA (Phyllanthus niruri Linn.) WITH PERMANGANOMETRY METHOD

*by Ai Sri Kosnayani*

---

**Submission date:** 19-Aug-2021 09:15PM (UTC+0700)

**Submission ID:** 1633230439

**File name:** RAN\_TEA\_Phyllanthus\_niruri\_Linn.\_WITH\_PERMANGANOMETRY\_METHOD.pdf (2.13M)

**Word count:** 1793

**Character count:** 9676

## A Qualitative Analysis of Tannin Type and Tannin Content in Meniran Tea (*Phyllanthus Niruri* Linn.) with Permanganometry Method

1<sup>st</sup> Ai Sri Kosnayani  
Department of Nutrition  
Siliwangi University, Indonesia  
aisrikosnayani@unsil.ac.id

2<sup>nd</sup> R. Reza El Akbar  
Department of Informatics Engineering  
Siliwangi University, Indonesia  
Reza@unsil.ac.id

3<sup>rd</sup> Liah Badriah  
Department of Biology Education  
Siliwangi University, Indonesia  
Liahbadriah014@gmail.com

4<sup>th</sup> Asep Kurnia Hidayat  
Department of Civil Engineering  
Siliwangi University, Indonesia  
asepkurnia@unsil.ac.id

**Abstract**—Meniran (*Phyllanthus niruri* Linn.) extract is known to be a source of antioxidants and non-toxic. The content of antioxidants in meniran is donated by tannin. The use of meniran (*Phyllanthus niruri* Linn.) extract as an herbal medicine has been widely used. Meniran (*Phyllanthus niruri* Linn.) can be brewed as tea and made into a drink, but the content of antioxidants in meniran tea is not widely known. The use of meniran (*Phyllanthus niruri* Linn.) tea can directly avoid the effects of residual solvents on extracting. The purpose of this study is to determine the type of tannin and tannin content in meniran (*Phyllanthus niruri* Linn.) tea with the permanganometry method. Meniran tea (*Phyllanthus niruri* Linn.) was made by washing and drying at room temperature. Meniran (*Phyllanthus niruri* Linn.) was collected from grass or roadside in the Tasikmalaya Regency. The research were conducted at the Basic Laboratory of the Faculty of Health Sciences, Siliwangi University. The results of the qualitative analysis showed that tannins contained in meniran tea are included in a group of hydrolyzed tannins with tannin content of catechol complex. The tannin content contained in 1 gram of meniran tea is 7.56 mg (7.56%).

**Keyword:** meniran, tannin, permanganometry, qualitative, antioxidants

### 1. INTRODUCTION

Meniran (*Phyllanthus niruri* Linn.) is an herbaceous plant that can live in tropical climate grasses. Meniran has been known as a drug in Asia, Africa and South America<sup>1</sup>. The Latin name meniran in India is *Phyllanthus amarus*<sup>2,3</sup>. Phytochemical analysis of meniran extract gave a positive reaction for groups of carbohydrates, alkaloids, flavonoids, glycosides, saponins, steroids and tannins<sup>4</sup> that provide antioxidant activity<sup>5</sup>.

Various studies on meniran extract showed that meniran extract was hypoglycemic, lowered cholesterol and triglycerides in alloxan-induced mice<sup>6</sup> and streptozotocin-induced<sup>7</sup>, hypotensive in male rabbits<sup>8</sup>. It can be used as the healing and / or prevention of infectious and degenerative diseases<sup>9</sup>, for improving insulin resistance in 10% sucrose-induced mice<sup>10</sup>, for providing anti-apoptotic effects and inhibiting inflammation<sup>11</sup>, and for reducing the weight of diabetic mice induced by alloxan<sup>12,13</sup>. In obese male rats, meniran water extracts can control weight gain, decrease blood glucose levels, increase HDL levels and reduce triglyceride levels<sup>14</sup>.

The flavonoid content in antioxidant meniran tea is partly contributed by tannin compounds. Tannin is a natural compound with a molecular weight of 500 - 3,000 with several free

hydroxyl phenol groups, forming stable bonds with proteins and biopolymers. The phenol groups found in tannins cause astringent, antiseptic effects that cause color if reacted with iron salts<sup>15</sup>

Meniran water extract (*Phyllanthus niruri* Linn.) can be produced in daily life by brewing directly with hot water and drink it like a tea. However, there is no research on the type of tannin and the levels of tannins in meniran tea. The importance of this research is to determine whether meniran tea is brewed directly containing tannins which have antioxidant activity for the body or not. If the meniran tea contains tannins with large amount then drinking meniran tea can be used as an option to consume medicinal plants.

## 2. MATERIALS AND METHOD

This research is an experimental research in the laboratory. The plant material used in this research was meniran tea (*Phyllanthus niruri* Linn.) taken from the area of Tasikmalaya Regency, with an antioxidant content of 74%<sup>14</sup>. Meniran tea used for each test was 2 grams. The chemicals used for this research were simplicia *Phyllanthus niruri* Linn powder, aquades,  $\text{KMnO}_4$  0.1N, indigo sulfonate acid LP, iron (III) ammonium sulfate or  $\text{FeCl}_3$ , ammonia Potassium iron (III) cyanide, hydrochloric acid, acetic acid, 10% Pb acetate, KBr 1% gelatin solution, Na acetate. Stiasny Formaldehyde.

The process were: weighed each  $\pm 2$  grams of meniran tea powder, given 50 ml of aquades, heated in water for 30 minutes then deposited and filtered with filter paper, and the filtrate was taken.

## 3. RESULTS AND DISCUSSION

### The Qualitative Analysis Results

The results obtained by reacting meniran tea with a solution of 1%  $\text{FeCl}_3$ , gelatin and a mixture of potassium ferisianida with ammonia, as shown as follows:



(a)



(b)



(c)

- (a) Change color to blue after addition of  $\text{FeCl}_3$  3% (positive tannin)
- (b) Formation of white deposits after the addition of 1% gelatin (positive tannin)
- (c) Change color to a collate after adding potassium cyanide and ammonia (positive tannin)

### Hydrolyzed tannin

Table 1. Results of hydrolyzed tannin type analysis test

Type test	Result
Meniran tea water plus $\text{FeCl}_3$	blue black color is formed
Meniran tea water plus HCl and then heated	red does not form
Meniran tea water plus 2 ml acetic acid and 1 ml Pb acetate solution 10%	formed sediment
Meniran tea water plus bromine (KBr) reagent	does not give sediment

### Condensed Tannin

Table 2 Results of condensed tannin type analysis test

Type test	Result
Meniran tea water plus $\text{FeCl}_3$ gives a greenish black color	Negative
Matchstick is forced into the meniran tea water, dried. Moistened with HCl and heated, phloroglucinol is formed which causes the matchstick to turn pink or red	Negative
Meniran tea water with HCl and heated, the red color of insolubela phlobagphenes	Negative
Meniran tea water plus 2 ml acetic acid and 1 ml Pb acetate solution 10%, do not cause sediment or remain in the form of solution	Negative
Meniran tea water plus bromine reagent (KBr), will give sediment	Negative

### Tannin Complex

Table 3 Results of complex tannin type analysis

Type test	Result
Meniran tea water is added with Stiasnyn L 3% formaldehyde - hydrochloric acid (2: 1) and refluxed for 30 minutes	Positive Red sediment
Meniran tea water plus $\text{FeCl}_3$	Negative greenish black color

### The Quantitative Analysis Result

Table 4 The results of quantitative analysis of 2 grams meniran tea with permanganometry

No sampel	Massa tannin (mg)	Tannin content
1	69	6,9%
2	80	8,0%
3	80	8,0%
4	80	8,0%
5	69	6,9%
Total	378	37,8
Average	75,6	7,56

### CONCLUSION

Meniran tea (*Phyllanthus niruri* Linn.) made by drying and brewing with hot water (without extraction) was positive for tannins. The types of tannins contained in meniran tea were hydrolyzed tannins and complex tannins containing catechol and galat. The content of tannins in 2 grams of meniran tea was 7.56%.

Meniran tea can be used as an alternative drink that contains antioxidants with various benefits. Further research is needed to analyze the levels of antioxidants contained in 1 cup of meniran tea because of antioxidants and not only given by tannins.

### ACKNOWLEDGEMENT

Thank you to LP2M PMP Siliwangi University for funding this research and the Dean of FIK Siliwangi University who has given permission to use the Nutrition Department Basic laboratory.

### REFERENCE

1. Mellinger CG, Carbonero ER, Noleto GR, et al. Chemical and biological properties of an arabinogalactan from *Phyllanthus niruri*. *J Nat Prod*. 2005;68(10):1479-1483. doi:10.1021/np050129s
2. Manjrekar AP, Jisha V, Bag PP, et al. Effect of *Phyllanthus niruri* Linn. treatment on liver, kidney and testes in CCl<sub>4</sub>induced hepatotoxic rats. *Indian J Exp Biol*. 2008;46(7):514-520.
3. Swamy VH., Patil M, Thippeswamy A., et al. Protective role of phyllanthus niruri extract in doxorubicin-induced myocardial toxicity in rats. *Indian J Pharmacol*. 2011;43(1):31. doi:10.3109/00016347709162115
4. Okoli C, Ezike A, Akah P. Evaluation of antidiabetic potentials of *Phyllanthus niruri* in alloxan diabetic rats Evaluation of antiplasmodial natural products View project Natural Products View project. *African J Biotechnol* . 2010;9(2):248-259. doi:10.5897/AJB09.872
5. S. Thiangthuma, B. Dejaeghera, M. Goodarzia, C. Tistaerta, A.Y. Gordienc, N. Nguyen Hoaid, M. Chau Vand, J. Quetin-Leclercq, L. Suntornsukb YVH. Potentially antioxidant compounds indicated from *Mallotus* and *Phyllanthus* species fingerprints. *J Chromatogr*. 2012:114-121. <http://dx.doi.org/10.1016/j.jchromb.2012.04.031>.

6. Okoli CO, Obidike IC, Ezike AC, Akah PA, Salawu OA. Studies on the possible mechanisms of antidiabetic activity of extract of aerial parts of *Phyllanthus niruri*. *Pharm Biol.* 2011;49(3):248-255. doi:10.3109/13880209.2010.501456
7. Nwanjo H. Studies On The Effect Of Aqueous Extract Of *Phyllanthus Niruri* Leaf On Plasma Glucose Level And Some Hepatospecific Markers in Diabetic Wistar Rats. *Internet J Lab Med.* 2006;2(2):1-6.
8. Amaechina FC, Omogbai EK. Hypotensive effect of aqueous extract of the leaves of *Phyllanthus amarus* Schum and Thonn (Euphorbiaceae). *Acta Pol Pharm - Drug Res.* 2007;64(6):547-552.
9. Oyewo, Emmanuel Bukoye, Akanji, Musbau Adewumi and Adekunle AS. Immunomodulation Capabilities of Aqueous Leaf Extract of *Phyllanthus amarus* in male Wistar Rats. *Rep Opin.* 2012;4(1):22-38. <http://www.sciencepub.net/report>.
10. Adeneye AA. The leaf and seed aqueous extract of *Phyllanthus amarus* improves insulin resistance diabetes in experimental animal studies. *J Ethnopharmacol.* 2012;144(3):705-711. doi:10.1016/j.jep.2012.10.017
11. Kandhare AD, Ghosh P, Ghule AE, Zambare GN, Bodhankar SL. Protective effect of *Phyllanthus amarus* by modulation of endogenous biomarkers and DNA damage in acetic acid induced ulcerative colitis: Role of phyllanthin and hypophyllanthin. *Apollo Med.* 2013;10(1):87-97. doi:10.1016/j.apme.2013.01.006
12. Adeneye AA, Amole OO, Adeneye AK. Hypoglycemic and hypocholesterolemic activities of the aqueous leaf and seed extract of *Phyllanthus amarus* in mice. *Fitoterapia.* 2006;77(7-8):511-514. doi:10.1016/j.fitote.2006.05.030
13. Shetti AA, Sanakal RD, Kaliwal BB. Antidiabetic effect of ethanolic leaf extract of *Phyllanthus amarus* in alloxan induced diabetic mice. 2012;2(1):11-15.
14. Ai Sri Kosnayani. Effects of *Phyllanthus niruri* Linn., metformin and combination of the both to body weight, fasting blood glucose, trygliserides and HDL in obese rats. In: Purwokerto: Proceeding of The 1st University of Muhammadiyah Purwokerto-Pharmacy International Conference; 2015:9-17. file:///D:/Referensi Teh Meniran/Kosnayani 2015.pdf%0D.
15. Throughput H, Neill MJO, Lewis JA, et al. Trease and Evans Pharmacognosy. *Eff Br mindfulness Interv acute pain Exp An Exam Individ Differ.* 2015;1:1689-1699. doi:10.1017/CBO9781107415324.004

# A QUALITATIVE ANALYSIS OF TANNIN TYPE AND TANNIN CONTENT IN MENIRAN TEA (*Phyllanthus niruri* Linn.) WITH PERMANGANOMETRY METHOD

## ORIGINALITY REPORT

**25%**  
SIMILARITY INDEX

**20%**  
INTERNET SOURCES

**22%**  
PUBLICATIONS

**16%**  
STUDENT PAPERS

## PRIMARY SOURCES

- |          |   |           |
|----------|---|-----------|
| <b>1</b> | <a href="http://www.dovepress.com">www.dovepress.com</a><br>Internet Source   | <b>2%</b> |
| <b>2</b> | <a href="http://vinikarus.wordpress.com">vinikarus.wordpress.com</a><br>Internet Source   | <b>1%</b> |
| <b>3</b> | Navneet Kaur, Baljinder Kaur, Geetika Sirhindi. " Phytochemistry and Pharmacology of L.: A Review ", Phytotherapy Research, 2017<br>Publication   | <b>1%</b> |
| <b>4</b> | Saravanan Shanmugam, Parimelazhagan Thangaraj, Bruno dos Santos Lima, Gabriela G.G. Trindade et al. "Protective effects of flavonoid composition rich <i>P. subpeltata</i> Ortega. on indomethacin induced experimental ulcerative colitis in rat models of inflammatory bowel diseases", Journal of Ethnopharmacology, 2020<br>Publication | <b>1%</b> |
| <b>5</b> | <a href="http://mafiadoc.com">mafiadoc.com</a><br>Internet Source   | <b>1%</b> |



6	<a href="http://mjpms.in">mjpms.in</a> Internet Source	1 %
7	Caroline G. Mellinger, Elaine R. Carbonero, Guilhermina R. Noletto, Thales R. Cipriani et al. " Chemical and Biological Properties of an Arabinogalactan from ", Journal of Natural Products, 2005 Publication	1 %
8	R. Reza el Akbar, Rahmi Nur Shofa, Muhammad Ilham Paripurna, Supratman. "The Implementation of Naïve Bayes Algorithm for Classifying Tweets Containing Hate Speech with Political Motive", 2019 International Conference on Sustainable Engineering and Creative Computing (ICSECC), 2019 Publication	1 %
9	<a href="http://www.eurekaselect.com">www.eurekaselect.com</a> Internet Source	1 %
10	<a href="http://www.koreascience.or.kr">www.koreascience.or.kr</a> Internet Source	1 %
11	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1 %
12	<a href="http://hal.archives-ouvertes.fr">hal.archives-ouvertes.fr</a> Internet Source	1 %



13	Maria Francilene Souza Silva, Fátima de Cássia E. de Oliveira, Maria Claudia dos Santos Luciano, Guilherme J. Zocolo et al. "Biochemical and Pharmaceutic Properties of Genus Phyllanthus spp.: Focus on P. amarus Schum. & Thonn. and P. niruri L.", Current Traditional Medicine, 2021 Publication	1 %
14	<a href="http://www.sciencedirect.com">www.sciencedirect.com</a> Internet Source	1 %
15	<a href="http://chsr.aua.am">chsr.aua.am</a> Internet Source	1 %
16	Submitted to School of Business and Management ITB Student Paper	1 %
17	<a href="http://greenlifecyprus.com">greenlifecyprus.com</a> Internet Source	1 %
18	<a href="http://utsouthwestern.influent.utsystem.edu">utsouthwestern.influent.utsystem.edu</a> Internet Source	1 %
19	Submitted to Study Group Australia Student Paper	1 %
20	<a href="http://19january2021snapshot.epa.gov">19january2021snapshot.epa.gov</a> Internet Source	1 %
21	Adejuwon Adewale Adeneye. "The leaf and seed aqueous extract of Phyllanthus amarus improves insulin resistance diabetes in	<1 %

experimental animal studies", Journal of  
Ethnopharmacology, 2012

Publication

---

22

Adeneye, A.A.. "Hypoglycemic and  
hypocholesterolemic activities of the aqueous  
leaf and seed extract of Phyllanthus amarus  
in mice", Fitoterapia, 200612

Publication

---

<1 %

23

Nundang Busaeri, Ida Ayu Dwi Giriantari,  
Wayan Gede Ariastina, IB Alit Swamardika.  
"Energy Management Assessment  
Measurement Using Three Measurement  
Approaches for Green Campus Concept",  
2019 International Conference on Sustainable  
Engineering and Creative Computing (ICSECC),  
2019

Publication

---

<1 %

24

Raja Noor Zaliha Raja Abd Rahman. "Anti-  
obesity Potential of Selected Tropical Plants  
via Pancreatic Lipase Inhibition", Advances in  
Obesity, Weight Management & Control, 2017

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off